

- Cauchy distribution, 59, 129–30, 807–8
 - and regression model, 285–86
 - translated, 129–30
- Cauchy-Schwartz inequality, 272
- Causality, 629–31
- Censored dependent variable models, 537–42
 - double censoring, 540–41
 - least squares estimation, 537–38
 - ML estimation, 538–40
 - tobit model, 537
- Censored sample, 534
- Centered sequence, 104
- Central limit theorem (CLT), 61, 62, 126–28, 134–35
 - condition CLT, 136
 - Lindeberg-Lévy, 135
 - Lyapunov, 135
 - McLeish, 135
 - multivariate, 136
 - simple, 126–27
- CG matrix, 252, 260–61
- Characterization of DGPs, complete and partial, 140
- Chebyshev inequality, 105, 799–800
- Chi-squared distribution, 808–9
 - central, 808
 - noncentral, 412–13, 449, 808–9
 - quadratic forms in normal vectors, 809
- Chow test, 375–81
 - assumption of homoskedasticity, 377
 - classic treatment, 375–76
 - heteroskedasticity-robust, 377
 - subsample too small, 378
 - and tests based on the GNR, 379
- Classical test statistics, 274–78, 435–57
 - asymptotic distribution under drifting DGP, 448–49
 - effect of reparametrization, 463–71
 - geometrical interpretation, 439–45
 - geometry of asymptotic distribution, 449–52
 - inequalities among, 456–57
 - for linear regression models, 452–57
 - numerical equality when Hessian is quadratic, 438–39
- Closeness (for vectors and matrices), 102
- Cochrane-Orcutt procedure, 334–35
 - estimated covariance matrix, 338–40
- Codimension (of a Euclidean subspace), 6
- Codomain (of a mapping), 777
- Coefficient of determination (R^2), 14–16
- Cofactor (for calculating determinant), 785
- Cointegrating vector, 716
 - asymptotically efficient estimation, 719
 - ML estimation, 726–29
 - OLS estimation, 717–19
 - super-consistent estimates, 718–19
- Cointegration, 715–19
 - bivariate example, 717
- Cointegration tests, 720–22
 - augmented Engle-Granger test, 720–21
 - Engle-Granger test, 720–21
 - LR tests in a VAR, 729–30
 - residual-based, 720–21
- Collinearity, 51, 181–86, 673–74
- Column vector, 770–71
- Common factor restrictions, 364–69
 - determining number of, 366–68
 - reason for name, 365–66
 - tests of, 366–68
- Compound event, 794
- Computability (of an estimator), 246
- Concentrated loglikelihood function, 267–69
 - and information matrix, 268–69
 - for LIML estimation, 647–48
 - for linear simultaneous equations model, 639–40
 - for multivariate regression model, 316–17
 - for nonlinear regression model, 280–81
 - for regression model with AR(1) errors, 346–47
 - for regression model with MA(1) errors, 354–56
 - for VAR(p) model, 685
- Condition CLT, 136
- Condition number (of a matrix), 181–82
- Condition WULLN, 137
- Conditional expectation, 131–32
- Conditional logit model, 532–33
 - independence of irrelevant alternatives, 533
- Conditional mean, 42, 52
- Conditional moment conditions
 - and choice of instruments, 602–7
 - and GMM estimation, 603–7
- Conditional moment (CM) tests, 571–81
 - computation via OPG regression, 574–77
 - and GMM estimation, 619–20
 - information matrix test, 578–81
 - and LM tests, 573
 - one-degree-of-freedom case, 572
 - and t tests, 572–73
 - Tauchen's variant, 577
 - tests for excess kurtosis, 572
 - tests for skewness, 572
- Confidence interval, 71–77