

SUPPLEMENT A  
(Data & Bootstrap)

For

Exploring the International Linkages of the Euro Area: a  
Global VAR Analysis

By

by S. Dees, F. di Mauro, M.H. Pesaran and L.V.Smith.

December 2006 (revised)

Part I: Data Sources and the Interpolation Procedure  
Part II. Bootstrapping the GVAR and Tests for Structural Stability

# 1 Data Sources and the Interpolation Procedure

The variables used in this paper are  $Y$ : Real GDP;  $CPI$ : Consumer price index;  $EQ$ : Equity price index;  $E$ : Exchange rate;  $R^S$ : Short-term interest rate;  $R^L$ : Long-term interest rate; and  $P^o$ : Oil price index.

## A.1. Real GDP

The source of all 33 countries, except Singapore, is the IMF's International Financial Statistics (IFS) GDP series in 1995 constant prices, except Australia (2001/02), Norway (2001), United Kingdom (2000) and United States (2000). France, Germany, Italy, Japan, South Africa, Spain, Netherlands, Switzerland, Australia, New Zealand, Mexico, United Kingdom and United States are all from series br, and the remaining countries are from series bp. Where recent data were not available, the IFS series were completed with growth rates derived from series provided by Global Insight. The data source for Singapore was Datastream.

Where quarterly data were not available (i.e. for Argentina, Belgium, Brazil, Chile, China India, Indonesia, Malaysia, Mexico, Philippines, Saudi Arabia, Thailand and Turkey), quarterly series were interpolated linearly from the annual series (see A.7). Interpolated series were used only during the periods 1981-1992 for Argentina, 1979 for Belgium, 1979-1989 for Brazil, 1979 for Chile, 1979-1996 for India, 1979-1982 for Indonesia, 1979-1987 for Malaysia, 1979 for Mexico, 1979-1980 for the Philippines, 1979-1992 for Thailand and 1979-1986 for Turkey. Quarterly output series were available for the subsequent periods.

For the period before the German unification, in 1990Q4, West German growth rates were used.

The data for Argentina, Austria, Belgium, Brazil, Chile, Finland, India, Indonesia, Korea, Malaysia, Mexico, Norway, Peru, Philippines, Sweden, Thailand and Turkey were seasonally adjusted. Seasonal adjustment was performed with EViews, using the U.S. Census Bureau's X12 program (for further details, see U.S. Census Bureau, 2000).

## A.2. Consumer Price Indices

The data source for all countries was the IFS Consumer Price Index series 64 zf, except China (64 xzf). The Price Index for China was seasonally adjusted post-1986.

## A.3. Equity Price Indices

The data source was the IFS series 62 zf (Industrial share prices) for 25 countries (Australia, Austria, Brazil, Canada, Chile, Finland, France, Germany, Italy, Japan, Korea, Mexico, Netherlands, New Zealand, Norway, Peru, Philippines, Saudi Arabia, South Africa, Spain, Sweden, Switzerland,

Thailand, United Kingdom, United States). For Norway, Sweden, France, United Kingdom, the IFS data were completed with OECD Main Economic Indicators database (MEI) the IFS data for Austria were completed with Datastream series.

The data source for Argentina, Belgium, Brazil and Switzerland was Datastream.

The data source for Malaysia, Turkey and China was Bloomberg.

#### A.4. Exchange Rates

IFS series rf was used for all countries.

#### A.5. Short-Term Interest Rates

The data source was the IFS series 60 b (Money market - interbank - rate) for 16 countries. For the 8 Euro Area countries (Austria, Belgium, France, Finland, Germany, Italy, Netherlands, and Spain) the ST interest rate was constructed as follows: for 1979Q1-1998Q4, the short-term country-specific inter-bank rate from IFS was used. From 99Q1-01Q4, the overnight EONIA rate was used as the common short-term interest rate for all 8 countries. IFS deposit rate series 60l were used for Argentina, Chile, Saudi Arabia and Turkey. The IFS Treasury Bill rate series 60c were used for Mexico and Philippines. For Sweden, some missing values at the end of the sample were replaced by the series 60 a (the two series are similar over the past). For China, New Zealand and Peru, IFS discount rate 60 were used. For India, Global Insight data were used to complete IFS series.

#### A.6. Long-Term Interest Rates

A long-term government bond rate was available from the IFS (series 61 zf) for 23 countries. Data from OECD were used to complete gaps in the IFS series for Austria and Sweden. Long-term interest rate series were not available for Argentina, Brazil, Chile, China, India, Indonesia, Peru, Saudi Arabia and Turkey.

#### A.7. Oil Price Index

For oil prices we used monthly averages of Brent Crude series from Datastream.

## 1.1 Description of the Interpolation Procedure

Let  $X_t$ ,  $t = 0, 1, 2, \dots, T$ , be the annual observations compiled as averages of  $m$  time-disaggregated observations,  $x_{it}$ ,  $i = 1, 2, \dots, m$ ,  $t = 1, 2, \dots, T$ , such that

$$X_t = \sum_{i=1}^m x_{it} \quad (1)$$

The objective is to estimate a relatively smooth set of observations,  $x_{ti}$ ,  $i = 1, 2, \dots, m$  that satisfy the above constraint. We confine ourselves to pure interpolation methods (namely without using any related economic time series) and assume that the underlying disaggregated observations are generated by the following time-varying first-order autoregressive process.

For year  $t$

$$\begin{aligned} x_{t1} &= \rho_t x_{t-1,m} + \mu_t \\ x_{t2} &= \rho_t x_{t1} + \mu_t \\ x_{t3} &= \rho_t x_{t2} + \mu_t \\ &\vdots \\ x_{tm} &= \rho_t x_{t,m-1} + \mu_t \end{aligned}$$

for year  $t+1$

$$\begin{aligned} x_{t+1,1} &= \rho_{t+1} x_{tm} + \mu_{t+1} \\ x_{t+1,2} &= \rho_{t+1} x_{t+1,1} + \mu_{t+1} \\ x_{t+1,3} &= \rho_{t+1} x_{t+1,2} + \mu_{t+1} \\ &\vdots \\ x_{t+1,m} &= \rho_{t+1} x_{t+1,m-1} + \mu_{t+1} \end{aligned}$$

and so on.

The above interpolations depend on the intercept coefficient,  $\mu_{t+1}$ , which is assumed to vary from one year to the next, and the autoregressive coefficient  $\rho_{t+1}$ . Solving for  $x_{t+1,i}$  recursively forward we have

$$x_{t+1,i} = \rho_{t+1}^i x_{tm} + \mu_{t+1} \frac{(1 - \rho_{t+1}^i)}{(1 - \rho_{t+1})}, \text{ for } i = 1, 2, \dots, m. \quad (2)$$

Substituting these in the constraint (1) we find

$$X_{t+1} = \rho_{t+1} \frac{(1 - \rho_{t+1}^m)}{(1 - \rho_{t+1})} x_{tm} + \frac{m\mu_{t+1}}{1 - \rho_{t+1}} - \rho_{t+1} \frac{(1 - \rho_{t+1}^m)}{(1 - \rho_{t+1})^2} \mu_{t+1}.$$

It is easily verified that the interpolations,  $x_{t+1,i}$ , do in fact exactly add up to the annual data,  $X_{t+1}$ .

The uniformly distributed interpolated series,  $x_{t+1,i} = X_{t+1}/m$ , for  $i = 1, 2, \dots, m$ , correspond to the case where  $\rho_{t+1} = 0$ . We adopt the geometrically (exponentially) interpolated series which is obtained by setting

$\mu_{t+1} = 0$ , while other intermediate cases can also be entertained, but in the case of our applications they tend to generate very similar outcomes.

For the exponential interpolation,  $\rho_{t+1}$  is computed as the solution to

$$X_{t+1} = \rho_{t+1} \frac{(1 - \rho_{t+1}^m)}{(1 - \rho_{t+1})} x_{tm}, \quad (3)$$

where  $x_{tm}$  is the observation at the end of the previous year. This formulation is suitable when interpolating the level of the variables (indices) rather than the growth rates and is applicable to I(1) variables.

To solve for  $\rho_{t+1}$ , let  $\lambda_{t+1,m} = X_{t+1}/x_{tm}$ , and write (3) in the expanded form

$$\rho_{t+1}^m + \rho_{t+1}^{m-1} + \dots + \rho_{t+1} = \lambda_{t+1,m}, \text{ for } t = 0, 1, \dots, \quad (4)$$

with

$$\lambda_{1,m} = X_1/x_{0m} = m(X_1/X_0). \quad (5)$$

It follows that

$$x_{t+1,i} = x_{tm} \rho_{t+1}^i, \quad t = 0, 1, \dots; i = 1, 2, \dots, m. \quad (6)$$

To proceed it is required to solve the  $m^{\text{th}}$  order polynomial equation given by (4). For the purpose of our empirical application we are interested in interpolating quarterly observations from annual series, which implies solving the quartic equation (for  $m = 4$ )

$$\rho_{t+1}^4 + \rho_{t+1}^3 + \dots + \rho_{t+1} - \lambda_{t+1,4} = 0. \quad (7)$$

To solve the quartic equation of the general form

$$A_4 z^4 + A_3 z^3 + A_2 z^2 + A_1 z + A_0 = 0$$

or

$$z^4 + a_3 z^3 + a_2 z^2 + a_1 z + a_0 = 0 \quad (8)$$

with  $a_i = A_i/A_4$ ,  $i = 0, 1, 2, 3$ , we substitute  $z = x - a_3/4$  in (8) which yields

$$x^4 + px^2 + qx + r = 0, \quad (9)$$

where

$$p = a_2 - \frac{3}{8}a_3^2, \quad q = a_1 - \frac{1}{2}a_2a_3 + \frac{1}{8}a_3^3$$

$$r = a_0 - \frac{1}{4}a_1a_3 + \frac{1}{16}a_2a_3^2 - \frac{3}{256}a_3^4.$$

In order to solve equation (9) it needs to be made factorable, which leads to the solution of the following cubic equation

$$u^3 + b_2u^2 + b_1u + b_0 = 0, \quad (10)$$

where

$$b_2 = -p, b_1 = -4r, b_0 = 4pr - q^2.$$

The cubic equation (10) has only one real root if the discriminant  $D$  is greater than zero, where  $D$  is defined by

$$D = Q^3 + R^2$$

and

$$Q = \frac{3b_1 - b_2^2}{9}, R = \frac{9b_1b_2 - 27b_0 - 2b_2^3}{54}.$$

In this case,  $D > 0$ , the unique real root is given by

$$u_1 = (R + \sqrt{D})^{1/3} - \frac{Q}{(R + \sqrt{D})^{1/3}} - \frac{1}{3}.$$

Then, by using the above solution to the cubic polynomial,  $u_1$ , the following quadratic equations arise

$$x^2 + \sqrt{u_1 - px} + \frac{1}{2}u_1 - \frac{q}{2\sqrt{u_1 - px}} = 0 \quad (11a)$$

$$x^2 - \sqrt{u_1 - px} + \frac{1}{2}u_1 + \frac{q}{2\sqrt{u_1 - px}} = 0, \quad (11b)$$

If  $x_r$  is a real solution of the pair of quadratics (11) then  $x_r - a_3/4$  is a real solution to the quartic equation (8). Thus a real solution to (7) is given by

$$\rho_{t+1} = x_{r,t+1} - 1/4$$

However, multiple real solutions can arise from the solution of the quartic equation defined by (8).

Consider two real solutions of (8),  $a$  and  $b$ . Let  $\{y_{i1}^a, y_{i2}^a, y_{i3}^a, y_{i4}^a, \dots\}$  and  $\{y_{i1}^b, y_{i2}^b, y_{i3}^b, y_{i4}^b, \dots\}$  be the levels of the interpolated series based on the choice of the roots  $a$  and  $b$ , respectively. In this case, we define

$$\Delta_a = \frac{|\ln(y_{i1}^a/y_{i1-1,4}^a)| + |\ln(y_{i21}^a/y_{i1}^a)| + |\ln(y_{i41}^a/y_{i3}^a)| + \dots}{4}$$

$$\Delta_b = \frac{|\ln(y_{i1}^b/y_{i1-1,4}^b)| + |\ln(y_{i21}^b/y_{i1}^b)| + |\ln(y_{i41}^b/y_{i3}^b)| + \dots}{4}$$

and choose  $a$  if  $\Delta_a < \Delta_b$ ,  $b$  otherwise.

## 2 Bootstrapping the GVAR and Tests for Structural Stability

To derive the empirical distribution of the structural stability tests and impulse response functions we employ the sieve bootstrap. The sieve bootstrap has been studied by Kreiss (1992), Bühlmann (1997) and Bickel and Bühlmann (1999) among others and has now become a standard tool when bootstrapping time series models.<sup>1</sup> The method rests on the assumption that the precise form of the parametric model generating the data is not known and that the true model belongs to the class of linear processes having an autoregressive representation of infinite order. Taking the estimated finite order vector autoregressive process that describes in our case the GVAR model to be an approximation to the underlying infinite order vector autoregressive process, we can use the sieve bootstrap for the basis of deriving critical values for the structural stability tests and for constructing bootstrap confidence regions.

In the case of stationary multivariate models, the sieve bootstrap has been used successfully to handle parameter estimation (Paparoditis, 1996). In the context of non-stationary time series, Park (2002) established an invariance principle applicable for the asymptotic analysis of the sieve bootstrap, which led Chang and Park (2003) to establish its asymptotic validity in the case of ADF unit root tests. Subsequently, Chang, Park and Song (2005) established the consistency of the sieve bootstrap for the OLS estimates of the cointegrating parameters assuming there exists one cointegrating relation amongst the variables under consideration. In what follows we consider the sieve bootstrap approach by resampling the residuals of the finite order global vector autoregressive process.

When bootstrapping unit root tests based on first order autoregressions, Basawa et al. (1991) show that the bootstrap samples need to be generated with the unit root imposed in order to achieve consistency for the bootstrap unit root tests. While our focus is not on bootstrapping unit root or cointegration tests, it seems natural to impose the unit root and cointegrating properties of the model when bootstrapping the statistics of interest. See also Li and Maddala (1997) who study the bootstrap cointegrating regression by means of simulation.

We begin by estimating the individual country VARX\*( $\hat{p}_i, \hat{q}_i$ ) models

---

<sup>1</sup>Another popular method is the block bootstrap by Künsch (1989). Choi and Hall (2000) discuss the substantial advantages of the sieve bootstrap over the block bootstrap for linear time series.

in their error correction form subject to reduced rank restrictions, for  $i = 0, 1, 2, \dots, N$  and  $t = 1, 2, \dots, T$ , where  $\hat{p}_i$  and  $\hat{q}_i$  are the estimated lag orders of the endogenous and foreign variables respectively based on the AIC. The estimated VARX $^*(\hat{p}_i, \hat{q}_i)$  are given by

$$\begin{aligned} \mathbf{x}_{it} = & \hat{\mathbf{a}}_{i0} + \hat{\mathbf{a}}_{i1}t + \hat{\mathbf{\Phi}}_{i1}\mathbf{x}_{i,t-1} + \dots + \hat{\mathbf{\Phi}}_{i\hat{p}_i}\mathbf{x}_{i,t-\hat{p}_i} \\ & + \hat{\mathbf{\Psi}}_{i0}\mathbf{x}_{it}^* + \hat{\mathbf{\Psi}}_{i1}\mathbf{x}_{i,t-1}^* + \dots + \hat{\mathbf{\Psi}}_{i\hat{q}_i}\mathbf{x}_{i,t-\hat{q}_i}^* + \hat{\mathbf{u}}_{it}, \end{aligned} \quad (12)$$

where we denote by  $\hat{r}_i$  the estimated number of cointegrating relations for country  $i$ . In estimating the cointegrating rank we entertain the case of an unrestricted intercept and restricted trend, the latter restricted to lie in the cointegrating space so as to avoid giving rise to quadratic trends in the level of the process.

The country specific models (12) are then combined via the link matrices  $W_i$  as described in Section 2, giving rise to the GVAR( $\hat{p}$ ) model expressed in terms of the global variables vector  $\mathbf{x}_t$  as

$$\hat{\mathbf{G}}\mathbf{x}_t = \hat{\mathbf{a}}_0 + \hat{\mathbf{a}}_1t + \hat{\mathbf{H}}_1\mathbf{x}_{t-1} + \dots + \hat{\mathbf{H}}_{\hat{p}}\mathbf{x}_{t-\hat{p}} + \hat{\mathbf{u}}_t \quad (13)$$

with  $\hat{p} = \max(\hat{p}_i, \hat{q}_i)$ , or alternatively,

$$\mathbf{x}_t = \hat{\mathbf{b}}_0 + \hat{\mathbf{b}}_1t + \hat{\mathbf{F}}_1\mathbf{x}_{t-1} + \dots + \hat{\mathbf{F}}_{\hat{p}}\mathbf{x}_{t-\hat{p}} + \hat{\varepsilon}_t \quad (14)$$

where  $\hat{\mathbf{F}}_j = \hat{\mathbf{G}}^{-1}\hat{\mathbf{H}}_j$ ,  $\hat{\mathbf{b}}_j = \hat{\mathbf{G}}^{-1}\hat{\mathbf{a}}_j$ , for  $j = 0, 1, \dots, \hat{p}$ ,  $\hat{\varepsilon}_t = \hat{\mathbf{G}}^{-1}\hat{\mathbf{u}}_t$  and  $\hat{\Sigma}_{\varepsilon} = \sum_{t=1}^T \hat{\varepsilon}_t\hat{\varepsilon}_t'/T$ . The total number of variables in the model is given by  $k = \sum_{i=0}^N k_i$  where  $k_i$  is the number of endogenous regressors in country  $i$ ,  $i = 0, 1, \dots, N$ .

Using the estimates from the fitted model (14) obtained from the observed data for  $\hat{p} = 2$ , we generate  $B$  bootstrap samples denoted by  $\mathbf{x}_t^{(b)}$ ,  $b = 1, \dots, B$ , from the process

$$\mathbf{x}_t^{(b)} = \hat{\mathbf{b}}_0 + \hat{\mathbf{b}}_1t + \hat{\mathbf{F}}_1\mathbf{x}_{t-1}^{(b)} + \hat{\mathbf{F}}_2\mathbf{x}_{t-2}^{(b)} + \varepsilon_t^{(b)}, \quad t = 1, 2, \dots, T, \quad (15)$$

by resampling the residuals  $\hat{\varepsilon}_t$  of the fitted model, with  $\mathbf{x}_0^{(b)} = \mathbf{x}_0$ , and  $\mathbf{x}_{-1}^{(b)} = \mathbf{x}_{-1}$ , where  $\mathbf{x}_0$  and  $\mathbf{x}_{-1}$  are the the actual initial data vectors. Prior to any resampling the residuals  $\hat{\varepsilon}_t$  are recentered to ensure that their bootstrap population mean is zero. The sieve bootstrap effectively reinterprets the familiar parametric AR model as a device for nonparametric estimation. The errors  $\varepsilon_t^{(b)}$  could also be drawn by parametric methods. Both these methods will be described in what follows. Simulating the GVAR model is clearly



preferable to simulating the country specific models separately. The latter requires that the country specific foreign variables,  $\mathbf{x}_{it}^*$ , and their lagged values are treated as strictly exogenous which might not be appropriate and could lead to unstable outcomes for  $\mathbf{x}_t$ .

It should be noted that the GVAR model given in (15) contains among others the inflation and real exchange rate variables. The choice of these variables rather than the price level and the nominal exchange rate was dictated by the results of the unit root tests. Once a set of  $\mathbf{x}_t^{(b)}$ ,  $b = 1, 2, \dots, B$  are generated, the price level and the nominal exchange rate are easily recovered, the foreign star variables are constructed using the weights in Table 2 and the inflation and real exchange rate variables are recreated.<sup>2</sup> Alternatively, the foreign (star) counterparts of these variables could be directly constructed, although it should be noted that this option is only valid in the case of fixed weights.

For each replication  $b$ , the individual country models are estimated in their error correction form, where  $p_i$ ,  $q_i$  and the number of the cointegrating relations,  $r_i$ , are fixed over all replications at the estimated values  $\hat{p}_i$ ,  $\hat{q}_i$  and  $\hat{r}_i$  obtained from the observed data, and a new set of VARX\* estimates are computed from

$$\begin{aligned} \mathbf{x}_{it}^{(b)} = & \hat{\mathbf{a}}_{i0}^{(b)} + \hat{\mathbf{a}}_{i1}^{(b)} t + \hat{\Phi}_{i1}^{(b)} \mathbf{x}_{i,t-1}^{(b)} + \dots + \hat{\Phi}_{i\hat{p}_i}^{(b)} \mathbf{x}_{i,t-\hat{p}_i}^{(b)} \\ & + \hat{\Psi}_{i0}^{(b)} \mathbf{x}_{it}^{*(b)} + \hat{\Psi}_{i1}^{(b)} \mathbf{x}_{i,t-1}^{*(b)} + \dots + \hat{\Psi}_{i\hat{q}_i}^{(b)} \mathbf{x}_{i,t-\hat{q}_i}^{*(b)} + \hat{\mathbf{u}}_{it}^{(b)}. \end{aligned} \quad (16)$$

We denote by  $ECM_{ij,t-1}^{(r)}$  the estimated error correction terms that correspond to the  $\hat{r}_i$  cointegrating relationships for country  $i$ , where  $i = 0, 1, \dots, N$  and  $j = 1, 2, \dots, \hat{r}_i$ .

## 2.1 Structural Stability Tests

For the structural stability tests consider the  $\ell^{th}$  equation of the estimated  $i^{th}$  country error correction model given by

$$\Delta x_{it,\ell} = \hat{\mu}_{i\ell} + \sum_{j=1}^{\hat{r}_i} \hat{\gamma}_{ij,\ell} ECM_{ij,t-1} + \sum_{n=1}^{\hat{p}_i} \hat{\phi}'_{in,\ell} \Delta \mathbf{x}_{i,t-n} + \sum_{s=0}^{\hat{q}_i} \hat{\psi}'_{is,\ell} \Delta \mathbf{x}_{i,t-s}^* + e_{it,\ell}, \quad (17)$$

which can be written more compactly as

---

<sup>2</sup>As the maximum order of the GVAR model is 2 in the current application, the actual data is used for the first two observations following such a transformation.

$$y_{it,\ell} = \hat{\theta}'_{it,\ell} \mathbf{z}_{it} + e_{it,\ell}, \quad (18)$$

where  $y_{it,\ell} = \Delta x_{it,\ell}$ ,  $\mathbf{z}_{it} = (1, E\hat{C}M'_{ij,t-1}, \Delta \mathbf{x}'_{i,t-n}, \Delta \mathbf{x}'_{i,t-s})'$  for  $j = 1, \dots, \hat{r}_i$ ,  $n = 1, \dots, \hat{p}_i$  and  $s = 0, \dots, \hat{q}_i$  and  $\hat{\theta}_{it,\ell} = (\hat{\mu}_{i\ell}, \hat{\gamma}_{ij,\ell}, \hat{\varphi}'_{in,\ell}, \hat{\vartheta}'_{is,\ell})'$ . Let  $e_{it,\ell}$  be the residuals from the estimated model (18) and  $\hat{\sigma}_{i\ell}^2 = T^{-1} \sum_{t=1}^T e_{it,\ell}^2$  the corresponding estimated error variance.

We consider a number of structural stability tests similar to those considered by Stock and Watson (1996). The null hypothesis for all the tests is that of parameter constancy, that is  $\theta_{\ell t} = \theta_{\ell}$ . The alternative varies depending on the test from non-stationarity, e.g random walks, to a one time change at an unknown break point for the sequential Wald type statistics, or some systematic movement in the parameters which we consider all to be subject to change. For expositional purposes we abstract from the index  $i$ .

### 2.1.1 1. Tests Based on the Cumulative Sum of OLS Residuals

The maximal OLS CUSUM statistic proposed by Ploberger and Krämer (1992) is similar to Brown, Durbin and Evans' (1975) CUSUM statistic although it is computed using OLS rather than recursive residuals. The mean square version of this test is also considered. Let  $\zeta_T(\delta) = \hat{\sigma}_{\ell}^{-1} T^{-1/2} \sum_{s=1}^{[T\delta]} e_{\ell s}$ , where  $[\cdot]$  is the greatest integer function, then

$$PK_{\text{sup}} = \sup_{\delta \in [0,1]} |\zeta_T(\delta)| \quad (19)$$

$$PK_{\text{msq}} = \int_0^1 \zeta_T(\delta)^2 d\delta \quad (20)$$

### 2.1.2 2. Random Walk Alternatives

Nyblom (1989) specifies as the alternative that  $\theta_{\ell t}$  follows a random walk, that is,  $\theta_{\ell t} = \theta_{\ell,t-1} + \eta_{\ell t}$ , where  $\eta_{\ell t}$  is i.i.d. and uncorrelated with error term corresponding to equation (18) and proposed the following statistic

$$\mathfrak{N}_{\ell} = T^{-2} \sum_{t=1}^T S'_{\ell t} \hat{\mathbf{V}}_{\ell}^{-1} S_{\ell t}, \quad (21)$$

where  $S_{\ell t} = \sum_{s=1}^t \mathbf{z}_s e_{\ell s}$  and  $\hat{\mathbf{V}}_{\ell} = (T^{-1} \sum_{t=1}^T \mathbf{z}_t \mathbf{z}'_t) \hat{\sigma}_{\ell}^2$ . The heteroskedasticity-robust version of the  $\mathfrak{N}_{\ell}$  statistic is obtained by replacing  $\hat{\mathbf{V}}_{\ell}$  in (21) with  $\tilde{\mathbf{V}}_{\ell} = T^{-1} \sum_{t=1}^T e_{\ell t}^2 \mathbf{z}_t \mathbf{z}'_t$ .

### 2.1.3 3. Sequential Wald Statistics

(i) Quandt (1960) likelihood ratio (QLR) statistic, in Wald form

$$QLR = \sup_{\delta \in (\delta_0, \delta_1)} F_{\ell T}(\delta)$$

(ii) The mean Wald statistic (Hansen (1992), Andrews and Ploberger (1994))

$$MW = \int_{\delta_0}^{\delta_1} F_{\ell T}(\delta) d\delta$$

(iii) The exponential average Wald statistic by Andrews and Ploberger (1994)

$$APW = \ln \left\{ \int_{\delta_0}^{\delta_1} \exp(F_{\ell T}(\delta)/2) d\delta \right\}.$$

To obtain the Wald statistic  $F_{\ell T}(\delta)$  for a break at  $t = m$ , where  $\delta = m/T$  or  $m = [T\delta]$  in all the above tests, equation (18) is initially estimated under the null of no structural change and the resulting sum of squares are defined as  $\mathcal{R}_\ell = \mathbf{e}'_\ell \mathbf{e}_\ell$ , where  $\mathbf{e}_\ell = (e_{\ell 1}, e_{\ell 2}, \dots, e_{\ell T})'$ . The model with a one time break at  $t = m$  is given by

$$\text{Subsample 1: } y_{\ell t} = \theta'_{1\ell} \mathbf{z}_t + \varepsilon_{1\ell t}, \quad t = 1, 2, \dots, m \quad (22)$$

$$\text{Subsample 2: } y_{\ell t} = \theta'_{2\ell} \mathbf{z}_t + \varepsilon_{2\ell t}, \quad t = m + 1, \dots, T. \quad (23)$$

Let  $e_{1\ell t}$  and  $e_{2\ell t}$  be the residuals from the OLS estimation of (22) and (23) respectively. Define  $\mathcal{R}_{1\ell} = \mathbf{e}'_{1\ell} \mathbf{e}_{1\ell}$  and  $\mathcal{R}_{2\ell} = \mathbf{e}'_{2\ell} \mathbf{e}_{2\ell}$ .

Then,

$$F_{\ell T}(\delta) = (T - 2\kappa) \frac{\mathcal{R}_\ell - \mathcal{R}_{\ell 1} - \mathcal{R}_{\ell 2}}{\mathcal{R}_{\ell 1} + \mathcal{R}_{\ell 2}}$$

where  $\kappa$  is the dimension of  $\theta_{it,\ell}$ , and  $\delta \in [\delta_0, \delta_1]$  with  $\delta_1 = 1 - \delta_0$ . The value for  $\delta_0$  was set to 0.25 and was chosen based on the maximum number of regressors over the individual VARX\* models.

The heteroskedasticity-robust version of the sequential Wald tests is given by

$$F_T(\delta) = (\hat{\mathbf{b}}_1 - \hat{\mathbf{b}}_2)' \mathbf{Q}_\ell^{-1} (\hat{\mathbf{b}}_1 - \hat{\mathbf{b}}_2)$$

where

$$\begin{aligned}\hat{\mathbf{b}}_1 &= (\mathbf{Z}'_1 \mathbf{Z}_1)^{-1} \mathbf{Z}'_1 \mathbf{Y}_{1\ell} \\ \hat{\mathbf{b}}_2 &= (\mathbf{Z}'_2 \mathbf{Z}_2)^{-1} \mathbf{Z}'_2 \mathbf{Y}_{2\ell}\end{aligned}$$

and

$$\begin{aligned}\mathbf{Q}_\ell &= (\mathbf{Z}'_1 \mathbf{Z}_1)^{-1} \left( \sum_{t=1}^m e_{\ell t}^2 \mathbf{z}_t \mathbf{z}'_t \right) (\mathbf{Z}'_1 \mathbf{Z}_1)^{-1} + \\ &\quad (\mathbf{Z}'_2 \mathbf{Z}_2)^{-1} \left[ \left( \sum_{t=1}^T e_{\ell t}^2 \mathbf{z}_t \mathbf{z}'_t \right) - \left( \sum_{t=1}^m e_{\ell t}^2 \mathbf{z}_t \mathbf{z}'_t \right) \right] (\mathbf{Z}'_2 \mathbf{Z}_2)^{-1}\end{aligned}$$

with  $\mathbf{Z} = (\mathbf{z}'_1, \dots, \mathbf{z}'_T)'$  and  $\mathbf{Y}_\ell = (y_{\ell 1}, \dots, y_{\ell T})'$ . Subscripts 1 and 2 refer to equations (22) and (23), respectively.

For each replication  $b$ , we consider the  $\ell$ th equation of the country-specific error correction models of  $\mathbf{x}_{it}^{(b)}$  given in (24)

$$\Delta x_{it,\ell}^{(b)} = \mu_{i\ell}^{(b)} + \sum_{j=1}^{\hat{r}_i} \gamma_{ij,\ell}^{(b)} ECM_{ij,t-1}^{(b)} + \sum_{n=1}^{\hat{p}_i} \hat{\varphi}_{in,\ell}^{(b)} \Delta \mathbf{x}_{i,t-n}^{(b)} + \sum_{s=0}^{\hat{q}_i} \vartheta_{is,\ell}^{(b)} \Delta \mathbf{x}_{i,t-s}^{*(b)} + e_{it,\ell}^{(b)}, \quad (24)$$

where  $ECM_{ij,t-1}^{(b)}$ ,  $j = 1, 2, \dots, \hat{r}_i$  are the estimated error correction terms corresponding to the  $\hat{r}_i$  cointegrating relations found for the  $i^{th}$  country based on the observed data and  $\ell = 1, \dots, k_i$ , and compute the above statistics  $W_{\ell s}^{(b)}$ ,  $WW_{\ell s}^{(b)}$ ,  $PK_{\text{sup}}^{(b)}$ ,  $PK_{\text{msq}}^{(b)}$ ,  $\mathfrak{N}_\ell^{(b)}$ ,  $QLR^{(b)}$ ,  $MW^{(b)}$  and  $APW^{(b)}$ . These statistics are then sorted into ascending order and their value which exceeds 95% of the observed statistics represents the appropriate 95% critical value for the structural stability tests.

## 2.2 Bootstrapping of Impulse Response Functions

On the assumption that the error term  $\mathbf{u}_t$  associated with equation (13) has a multivariate normal distribution, the  $k \times 1$  vector of the generalized impulse response functions in the case of a one standard error shock to the  $j^{th}$  equation corresponding to a particular shock in a particular country on  $\mathbf{x}_{t+n}$  is given by

$$GI_{j,n} = \frac{\tilde{\mathbf{F}}^n \mathbf{G}^{-1} \Sigma_u \mathbf{e}_j}{\sqrt{\mathbf{e}'_j \Sigma_u \mathbf{e}_j}}, \quad n = 0, 1, 2, \dots \quad (25)$$

where  $\mathbf{e}_j$  is a  $k \times 1$  selection vector with unity as its  $j^{th}$  element,  $\Sigma_u$  is the covariance matrix of  $\mathbf{u}_t$ , and  $\tilde{\mathbf{F}} = \mathbf{E}_1 \mathbf{F} \mathbf{E}'_1$ , with  $\mathbf{F} = \begin{pmatrix} \mathbf{F}_1 & \mathbf{F}_2 \\ \mathbf{I}_k & \mathbf{0} \end{pmatrix}$  and

$\mathbf{E}_1 = \begin{pmatrix} \mathbf{I}_k & \mathbf{0}_{k \times k} \end{pmatrix}$ , which follows from re-writing (14) in its companion form.<sup>3</sup> This result also holds in non-Gaussian but linear settings where the conditional expectations can be assumed to be linear.

In the case of a structural shock the corresponding generalized impulse response function is given by

$$SI_{j,n} = \frac{\tilde{\mathbf{F}}^n(\mathbf{P}_G^0 \mathbf{G})^{-1} \boldsymbol{\Sigma}_u \boldsymbol{\epsilon}_j}{\sqrt{\boldsymbol{\epsilon}_j' \boldsymbol{\Sigma}_u \boldsymbol{\epsilon}_j}}, \quad n = 0, 1, 2, \dots \quad (26)$$

where  $\mathbf{P}_G^0$  is defined in Section 7.

For each bootstrap replication, having estimated the individual country models using the simulated data  $\mathbf{x}_t^{(b)}$ , the GVAR is reconstructed as described above and the impulse responses are calculated based on the formulas (25) and (26) as  $GI_{j,n}^{(b)}, SI_{j,n}^{(b)} \forall n$ . These statistics are then sorted into ascending order  $\forall n$  and the  $(1 - \gamma)100\%$  confidence interval is calculated by using the  $\gamma/2$  and  $(1 - \gamma/2)$  quantiles, say  $s_{\gamma/2}$  and  $s_{(1-\gamma/2)}$ , respectively of the bootstrap distribution of  $GI_{j,n}$  and  $SI_{j,n}$ .

## 2.3 Generating the Simulated Errors

### 2.3.1 Parametric Approach

Under the parametric approach the errors are generated from a multivariate distribution with zero means and covariance matrix  $\hat{\boldsymbol{\Sigma}}_\varepsilon$  given by  $\hat{\boldsymbol{\Sigma}}_\varepsilon = \frac{1}{T} \sum_{t=1}^T \hat{\boldsymbol{\varepsilon}}_t \hat{\boldsymbol{\varepsilon}}_t'$ . To obtain the simulated errors for the  $k$  variables in the GVAR model we first generate  $kT$  draws from an i.i.d distribution which we denote by  $\mathbf{v}_t^{(b)}$ ,  $t = 1, 2, \dots, T$ . In our application we generate  $\mathbf{v}_t^{(b)}$  as  $IIN(0, \mathbf{I}_k)$  although other parametric distributions could also be entertained. Invoking the spectral decomposition, the variance-covariance matrix of the estimated GVAR residuals are decomposed as  $\hat{\boldsymbol{\Sigma}}_\varepsilon = \hat{\mathbf{P}} \hat{\boldsymbol{\Lambda}} \hat{\mathbf{P}}'$ , where  $\hat{\boldsymbol{\Lambda}}$  is a diagonal matrix containing the eigenvalues of  $\hat{\boldsymbol{\Sigma}}_\varepsilon$  on its diagonal and  $\hat{\mathbf{P}}$  is an orthogonal matrix consisting of its eigenvectors. Note that the Choleski decomposition of  $\hat{\boldsymbol{\Sigma}}_\varepsilon$  is not applicable in this case due to the semi-positive definite nature of this matrix that follows from the underlying common factor structure of the GVAR. The errors  $\boldsymbol{\varepsilon}_t^{(b)}$ ,  $t = 1, 2, \dots, T$ , are then computed as  $\boldsymbol{\varepsilon}_t^{(b)} = \hat{\mathbf{A}} \mathbf{v}_t^{(b)}$ , where  $\hat{\mathbf{A}} = \hat{\mathbf{P}} \hat{\boldsymbol{\Lambda}}^{1/2}$ .

---

<sup>3</sup>See Pesaran and Shin (1998) for further details.

### 2.3.2 Non-Parametric Approach

To obtain a bootstrap sample for the  $k$  variables in the GVAR model, we initially pre-whiten the residuals  $\hat{\eta}_t$  by using the generalized inverse of  $\hat{\mathbf{A}}$  as given above, denoted  $\hat{\mathbf{A}}_g^-$ , so that  $\hat{\eta}_t = \hat{\mathbf{A}}_g^- \varepsilon_t$ . The generalized inverse of  $\hat{\mathbf{A}}$  is required due to the semi-positive definite nature of this matrix as was pointed out earlier. We then resample with replacement from the  $kT$  elements of the matrix obtained from stacking of the vectors  $\hat{\eta}_t$ , for  $t = 1, 2, \dots, T$ . This is done in order to reduce the repetition of the bootstrap samples. The bootstrap error vector is then obtained as  $\varepsilon_t^{(b)} = \hat{\mathbf{A}} \hat{\eta}_t^{(b)}$ , where  $\hat{\mathbf{A}}$  is the same as above, and  $\hat{\eta}_t^{(b)}$  is the  $k \times 1$  vector of re-sampled values from  $(\hat{\eta}_1, \hat{\eta}_2, \dots, \hat{\eta}_T)$ .

## References

- [1] Andrews, D.W.K. and Ploberger, W. (1994). Optimal Tests When a Nuisance Parameter is Present Only under the Alternative, *Econometrica*, 62, 1383-1414.
- [2] Basawa, I.V., Mallik, A.K., McCormick, W.P., Reeves, J.H. and Taylor, R.L. (1991). Bootstrapping Unstable First-Order Autoregressive Processes, *Annals of Statistics*, 1098-1101.
- [3] Bickel, P.J., and Bühlmann, P. (1999) A New Mixing Notion and Functional Central Limit Theorems for a Sieve Bootstrap in Time Series, *Bernoulli*, 5, 413-446.
- [4] Brown, R.L., Durbin, J. and Evans, J.M. (1975). Techniques for Testing the Constancy of Regression Relationships Over Time, *Journal of Royal Statistical Society, Series B*, 37, 149-163.
- [5] Bühlmann, P. (1997). Sieve Bootstrap for Time Series, *Bernoulli* 3, 123-148.
- [6] Chang, Y. and Park, J.Y. (2003). A Sieve Bootstrap for the Test of a Unit Root, *Journal of Time Series Analysis*, 24, 379-400.
- [7] Chang, Y., Park, J.Y. and Song, K. (2005). Bootstrapping Cointegrating Regressions, forthcoming in the *Journal of Econometrics*.
- [8] Choi, E. and Hall, P. (2000). Bootstrap Confidence Regions Computed From Autoregressions of Arbitrary Order, *Journal of the Royal Statistical Society, Series B*, 62, 461-477.

- [9] Hansen, B.E. (1992). Tests for Parameter Instability in Regressions with I(1) Processes, *Journal of Business and Economic Statistics*, 10, 321-336.
- [10] Kreiss, J.P. (1992). Bootstrap procedures for AR( $\infty$ )-processes. In: Jockel, K.H., Rothe, G., Sandler, W. (Eds.), *Bootstrapping and Related Techniques. Lecture Notes in Economics and Mathematical Systems*, Vol. 376. Heidelberg, Springer-Verlag, 107–113.
- [11] Künsch, H.R. (1989). The Jackknife and the Bootstrap for General Stationary Observations, *Annals of Statistics*, 17, 1217-1241.
- [12] Li, H. and Maddala, G.S. (1997). Bootstrapping Cointegrating Regressions, *Journal of Econometrics*, 80, 297-318.
- [13] Nyblom, J. (1989). Testing for the Constancy of Parameters Over Time. *Journal of the American Statistical Association*, 84, 223-230.
- [14] Paparoditis, E. (1996). Bootstrapping Autoregressive and Moving Average Parameter Estimates of Infinite Order Vector Autoregressive Processes. *Journal of Multivariate Analysis*, 57, 277–296.
- [15] Park, J.Y. (2002). An Invariance Principle for Sieve Bootstrap in Time Series, *Econometric Theory*, 18, 469-490.
- [16] Pesaran, M.H. and Shin, Y. (1998). Generalized Impulse Response Analysis in Linear Multivariate Models, *Economics Letters*, 58, 17-29.
- [17] Ploberger, W. and Krämer, W. (1992). The CUSUM test with OLS Residuals, *Econometrica*, 60, 271-286.
- [18] Quandt, R. (1960). Tests of the Hypothesis that a Linear Regression System Obeys two Separate Regimes, *Journal of the American Statistical Association*, 55, 324-330.
- [19] Stock, J.H. and Watson, M.W. (1996). Evidence on Structural Instability in Macroeconomic Time Series Relations, *Journal of Business and Economic Statistics*, 14, 11-30.