

---

**Addendum to NOTE 4****Sampling Distribution of OLS Estimator of  $\beta_2$** **A Monte Carlo Simulation**

***The True Model:*** is given by the **population regression equation (PRE)**

$$Y_i = \beta_1 + \beta_2 X_i + u_i = 90 + 0.90 X_i + u_i \quad (1)$$

where

$$\beta_1 = 90.0 \quad \text{and} \quad \beta_2 = 0.90;$$

$Y_i$  = weekly consumption expenditures of the  $i$ -th household;

$X_i$  = weekly disposable income of the  $i$ -th household;

$u_i$  = an iid random error term that is assumed to be  $N(0, \sigma^2)$ .

***Two Alternative Models:*** specify different values for  $\sigma^2 = \text{Var}(u_i | X_i)$ .

**Model 1:** sets  $\sigma^2 = \text{Var}(u_i | X_i) = 900$ ,  $\sigma = \sqrt{\text{Var}(u_i | X_i)} = \text{se}(u_i | X_i) = 30$ .

**Model 2:** sets  $\sigma^2 = \text{Var}(u_i | X_i) = 32,400$ ,  $\sigma = \sqrt{\text{Var}(u_i | X_i)} = \text{se}(u_i | X_i) = 180$ .

---

### *The Monte Carlo Simulations*

- Set  $N = 40$  observations.
- Set populations values of  $X$ ,  $\beta_1$  and  $\beta_2$ , and  $\sigma^2 = \text{Var}(u_i | X_i)$ .
- Generate 1,000 independent random samples of  $Y_i$ ,  $X_i$  and  $u_i$  values.
- For each of these 1,000 independent random samples, compute the values of the OLS coefficient estimators:

$$\hat{\beta}_2 = \frac{\sum_i x_i y_i}{\sum_i x_i^2} \quad (2)$$

$$\hat{\beta}_1 = \bar{Y} - \hat{\beta}_2 \bar{X} \quad (3)$$

where  $x_i \equiv X_i - \bar{X}$ ,  $y_i \equiv Y_i - \bar{Y}$ ,  $\bar{X} = \sum_i X_i / N$ , and  $\bar{Y} = \sum_i Y_i / N$ .

- Tabulate the 1,000 estimates of  $\beta_2$ , and the 1,000 estimates of  $\beta_1$ .

Summary statistics for simulation of Model 1 for which  $\sigma^2 = 900$ ,  $\sigma = 30$ :

```
. summarize
```

Variable	Obs	Mean	Std. Dev.	Min	Max
b1ols	1000	<b>90.12619</b>	10.07493	61.00864	129.009
b2ols	1000	<b>.8998154</b>	.008213	.8695259	.9269965


Summary statistics for simulation of Model 2 for which  $\sigma^2 = 32,400$ ,  $\sigma = 180$ :

```
. summarize
```

Variable	Obs	Mean	Std. Dev.	Min	Max
b1ols	1000	<b>90.75711</b>	60.44958	-83.94814	324.0542
b2ols	1000	<b>.8988922</b>	.0492779	.7171556	1.061979


- Comparison of Sampling Distributions of  $\hat{\beta}_2 = \frac{\sum_i x_i y_i}{\sum_i x_i^2}$  :

Sampling distribution of OLS estimator of beta2: beta2 = 0.90, N = 40



Model 1:  $E(u) = 0$ ,  $Var(u) = 900$ ,  $SE(u) = 30$ ,  $N = 40$

Sampling distribution of OLS estimator of beta2: beta2 = 0.90, N = 40



Model 2:  $E(u) = 0$ ,  $Var(u) = 32,400$ ,  $SE(u) = 180$ ,  $N = 40$